



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

all around, and the very broad central area occupied by irregular, more or less elongated reticulations. Thus the scale comes to closely resemble those of the Mormyridæ. As it now seems evident that the ancestors of the Teleosteans must have had reticulated scales, or at least that the ordinary radial sculpture is derived from the reticulated type, this *Moxostoma* scale must be regarded as uniquely primitive or atavistic for the general group to which it belongs.

3. Dr. G. A. Boulenger has very kindly sent me scales of the cæciliid amphibian *Ichthyophis glutinosus*. These are very small, embedded in the skin, cycloid in form. The pattern is extremely characteristic, consisting of concentric grooves connected at intervals by cross-lines, the whole effect being like that of bricks in a wall. The concentric grooves are probably not circuli, nor can I make out anything corresponding to the circuli of fishes. In parts of the scales, however, the markings become irregular, producing a reticulation which closely simulates that of the reticulate-scaled fishes. I believe that the scales are really comparable to fish-scales, and that the sculpture is the same as the radial sculpture of fishes. No fish scale has been seen resembling in detail that of *Ichthyophis*; such scales as those of *Chrosomus* are superficially similar, but owe their circular lines to different elements.¹

T. D. A. COCKERELL

UNIVERSITY OF COLORADO

NOTES ON THE GENUS *TYPHA* AND ITS NEMATODE
ROOT GALL—*HETERODERA RADICICOLA*
(GREEFF) MULL.

DURING the summer of 1908, while investigating some problems connected with the root system of *Typha latifolia*, I found a number of abnormal growths on the rootlets. These growths appeared as irregularly spherical or fusiform enlargements, varying in size from 1 to 5 mm. in diameter. They were identified by Professor Atkinson as root galls caused

¹ Since this was written, I have found that a deep-sea eel, *Synaphobranchus pinnatus*, has scales curiously similar to those of *Ichthyophis*.

by the nematode *Heterodera radiculicola*. I have collected these galls at the same station (limnology station of Cornell University) three successive years, but have never found them on *Typha* in any other locality.

Professor Atkinson¹ thought, from his observations of this worm on potatoes and tomatoes, that, if favorable opportunity should occur for its introduction in the north, it might become a pest. Webber and Orton² say it will never become a serious pest in the north, as severe cold kills the worm. Van Hook³ reports the worm as wintering in ginseng beds which had been mulched and also in protected forest beds. This worm has been a serious pest to ginseng in the north.

Stone and Smith⁴ found the galls on outdoor plants, but concluded that they were transient.

The plants observed by me in the Cayuga marshes are located along the shore line of one of the arms of Fall Creek where moisture is plentiful in the soil all winter. Winter observations prove that the soil in which the galls are found does not freeze. None of the galls have been found more than eighteen inches below the surface.

L. N. HAWKINS

CORRELATION NOTES

IN describing the fauna of the Moorefield shales of Arkansas¹ Mr. George H. Girty lists and describes the following fossils among others from the region: *Productus inflatus* var. *coloradoensis* Girty (?),² *Productus arkansanus* var. *multiliratus* Girty,³ and *Diaphragmus elegans* Norwood and Pratten.⁴ By a comparison of the figures of these fossils on plate iv.⁵ with fossils which the writer collected

¹ Bull. 9, Alabama Exp. Sta.

² U. S. Dept. Agr., Bur. Plant Ind., Bull. 17, 1902.

³ Cornell Agr. Exp. Sta., Bull. 219, 1904.

⁴ Bull. 55, Mass. Agr. Exp. Sta., 1898.

⁵ "The Fauna of the Moorefield Shale of Arkansas," U. S. Geol. Survey, Bulletin No. 439.

² *Ibid.*, pp. 42-43.

³ *Ibid.*, p. 43.

⁴ *Ibid.*, pp. 51-52.

⁵ *Ibid.*, plate iv.